



Find A Hospital Why VCA
In The Community
Know Your Pet

First Aid for Dogs

By Ernest Ward, DVM | Emergency Situations

Emergencies come in all forms; automobile accidents, bite wounds, burns, heatstroke, poisoning, seizures, and more.

For a general overview of what constitutes an emergency, and how to handle common crisis situations, refer to our fact sheet on Canine Emergencies.

What is first aid?

First aid is initial treatment given in a medical emergency. Its purpose is:

- To preserve life
- To reduce pain and discomfort
- To minimize any risk of permanent disability or disfigurement

In an emergency, what should I do first?

1. Keep calm and assess the scene for any additional threats to you or your pet. This is important for everyone's safety.
2. Keep your dog warm, as quiet as possible, and keep movement to a minimum, especially if there is possible trauma, broken limbs, or any neurological symptoms.

3. Contact your veterinary hospital, inform them of the situation and get specific first aid advice.
4. To safely move or transport an injured dog, get somebody to help you. For a small dog, put it into its carrier (remove the top for easy and safe access to the carrier; don't push an injured dog through the small door or opening), or use a suitable container such as a strong cardboard box. For a larger dog, use a makeshift stretcher made out of some rigid material such as an appropriate sized, sturdy piece of wood. Carefully maneuver the dog onto a blanket or coat so that it can be gently moved to the carrier, box or stretcher. The blanket will help stabilize the neck and spine, and prevent inadvertent biting or scratching from the injured pet.
5. Get to a veterinary hospital as soon as possible, but drive carefully!

What are some tips on restraining or calming an injured dog?

The majority of injured animals will be panicked and/or disoriented.

“The stress of an emergency involving a pet or owner can cause an otherwise friendly animal to act aggressively.”

The stress of an emergency involving a pet or owner can cause an otherwise friendly animal to act aggressively. Although most panicky dogs will respond to a calm, soothing voice and gentle stroking of the head and shoulders, use caution when approaching or touching any injured animal. It is important to ensure the safety of all rescue personnel that are attempting to assist with an injured animal. Some of the types of restraint that can ensure the safety of both dog and humans include:

- **Muzzling:** You can create a muzzle out of a leash, belt, sock, rope or strap. Loop the cord around the muzzle and tighten it to prevent the animal from biting. Dogs and cats have only one muscle to open their jaw so once the jaw is closed, it is relatively easy to hold it safely shut. Animals can breathe through their nostrils unless the nose is injured or obstructed.
- **Wrapping:** You can wrap the body of a fractious pet in a bag or towel. Be sure to keep the head exposed and not constrict the trachea.
- **Immobilizing:** If you are suspicious of spinal injury, lay the animal on a board and secure it on the board with straps or cords. Pay special attention to

immobilizing the head and neck.

What are the signs of shock?

Shock is a complex systemic or whole body reaction to a number of emergency situations. These include severe trauma, hemorrhage or sudden loss of blood, heart failure and other causes of decreased circulation (e.g. severe and sudden allergic reaction and heat stroke). A life-threatening fall in blood pressure is a dangerous part of shock.

“Systemic shock may cause irreversible injury to body cells, and it can be fatal.”

If not treated quickly and effectively, systemic shock may cause irreversible injury to body cells, and it can be fatal.

Clinical signs of systemic shock include rapid breathing and elevated heart rate with pale mucous membranes: gums, lips or under the eyelids. The feet or ears may feel cold and your dog may vomit and shiver. As shock progresses most pets become quiet and unresponsive.

What should I do if my dog is showing signs of shock?

Keep the dog as quiet as possible and try to conserve heat by covering it with blankets, towels, or even newspapers.

Follow the **A, B, C's** of first aid:

- A** **Airway**
- B** **Breathing**
- C** **Cardiac function**

Airway - Anything that obstructs the airway prevents oxygen entering the lungs. Do your best to clear the mouth and throat of any obstruction such as vomit, saliva or foreign bodies such as grass, sticks or balls. **Be careful**; your pet may bite you in panic.

“If you are unsure about the health or vaccination status of the injured pet, avoid contact with bodily fluids and blood.”

Breathing - If the dog is unconscious and does not appear to be breathing, try gently pumping the chest with the palm of your hand, at the same time feeling just behind the elbow to detect a heartbeat or pulse. If this is unsuccessful, give the pet rescue breathing (see below). Be careful! Injured pets may bite you out of fear. If you are unsure about the health or vaccination status of the injured pet, avoid contact with bodily fluids and blood.

Cardiac function - If you are unable to detect a heartbeat or pulse, or if appears weak and slow, try pressing on the chest with your palm and elevate the lower half of the body to promote blood flow to the brain. Follow the steps below, under CPR.

How do I perform Rescue Breathing for a dog?

When you encounter an unresponsive dog, the first step is to ensure that there is an open airway.

1. Carefully pull the tongue out of the mouth.
2. Extend the head and neck so that they are in a straight line. DO NOT overextend the neck in animals that have obvious head and neck trauma.
3. Carefully clear the mouth of any debris that may be obstructing breathing.
4. Place a resuscitator mask or your hand over the animal's muzzle while holding the mouth shut and neck extended. For small dogs, you can sometimes improvise with a styrofoam cup or other similar item, by placing the opening over the dog's face and poking a large hole in the bottom for you to breathe through. Ensure a relatively tight seal around the muzzle.
5. Blowing into the nostrils, give 2-3 breaths and watch for a rise in the chest. If you do not see a rise in the chest, reposition the neck or search for airway obstruction.
6. If you believe there is an airway obstruction that you cannot see, turn the dogs upside down, with the back against your chest. Give 5 sharp thrusts to the

abdomen to try and expel any object (this maneuver can be difficult to do in large dogs!).

7. For rescue breathing, provide 20 breaths per minute.

8. If the animal fails to breathe on its own, you may attempt an acupuncture maneuver. Press firmly with your fingernail or other sharp object in the space just beneath the nose on the upper lip (nasal philtrum). Maintain the pressure for 10-30 seconds.

What if the dog requires CPR?

After you have established an airway and begun rescue breathing, if there are still no obvious signs of life you should attempt cardiac compressions.

1. Make sure there is no major bleeding. If there is bleeding, have an assistant manage the bleeding (see below) while you perform CPR.
2. If possible, lay the dog on its right side.
3. Feel for a heartbeat or femoral pulse. The femoral pulse is located inside the leg in the groin region. Dogs DO NOT have a readily palpable carotid (neck) pulse.
4. Bend the left forearm and note the location where the elbow touches the chest. This is close to the middle of the rib cage.
5. Placing one hand on each side of the chest in the middle of the rib cage, vigorously compress the chest 3-5 times followed by 2 rescue breaths. The rate should be about 3 compressions every 2 seconds.
6. Try to compress the chest wall at least 30%. This is ½-1" (1-2 cm) in small dogs and 1½-2" (3-5 cm) in larger dogs.

What is the specific first aid for some of the more common emergencies?

Blood loss

Once you have followed A, B, C above, if the bleeding is still severe, try to stop it. If bleeding is from a cut pad or paw, apply a dressing using a piece of absorbent bandage or clothing. If the bleeding persists and is soaking through the bandage, don't waste any more time, since this is a medical emergency. Most bleeding

wounds will require medical or surgical treatment. If the wounds are treated within four (4) hours, they can often be sutured. Deep cuts treated after four hours have increased risk of infection and complication, and require more extensive surgery.



Burns and scalds

Cool the burned area with cold water as quickly as possible. Cover the burned area with damp towels. If the injury is due to a caustic substance, rinse with cold water for fifteen (15) minutes and contact your veterinarian for further advice. Animals that have been exposed to heat or smoke from a fire should be offered water as soon as the situation is stable.

Eye injuries

Injuries to the eye are always very painful and can threaten the eyesight. If a foreign body (grass awn, stick, etc.) can be seen, it may be possible to remove it by gently rinsing the eye with eyewash or contact lens saline solution. Do not allow the dog to rub the eye, either with its paws or against the furniture or carpet. Seek veterinary advice as soon as possible.

Seizures

Seizures can be due to many causes. These range from eclampsia (milk fever in a nursing mother) to epilepsy. If due to eclampsia, remove the puppies from the mother immediately. All dogs that are seizing or have had a recent seizure should be kept in a dark, quiet, confined area until medical help can be sought. Contact your veterinarian immediately.

Heat stroke

This most commonly occurs in hot weather when dogs are left in cars without adequate ventilation. Body temperature rises dramatically. Initial clinical signs include excessive panting and obvious distress, but can quickly progress to coma and death. Reduce the pet's body temperature as quickly as possible using cool water and keep the dog wet during transport to the veterinarian. Keep the car windows open. Evaporation will help reduce body temperature. Avoid using ice or ice water because this may drop the temperature too quickly and cause additional complications.

As a final note, **after being involved in an emergency or accident, it is important that you take your dog for a veterinary examination as soon as possible, even if it appears to have recovered fully.**

This client information sheet is based on material written by: Ernest Ward, DVM

PET FIRST AID KIT

Phone numbers: Your Veterinarian, nearest ER clinic with directions, Poison Control Center

Paperwork: Copy of Rabies Certificate and Photo (in case pet is lost)

Nylon leash

Self-cling bandage

Roll gauze

Gauze pads

Medical tape

Cotton balls and Q-tips

Disposable gloves

Rectal thermometer and lubricant (normal temp 100-102F)

Blunt end scissors

Small hemostats (vs. tweezers)

Nail clippers

Kwik stop

Small light source

Betadine scrub

3% hydrogen peroxide

Rubbing alcohol

Eye irrigation solution (exp.)

Triple antibiotic ointment (exp.)

Cortisone cream (exp.)


Diphenhydramine/Benadryl (1mg/pound orally) antihistamine, mild tranquilizer (exp.)

Gas X (exp.)


Famotidine/Pepcid (0.5mg/kg orally) (exp.)

Other medications based on your pet's health needs – consult your veterinarian

Vcahospitals.com/know-your-pet/common-emergencies-in-dogs



45° cut
removes tip
of nail.
Minimise
nail impact
on the floor.



Grind / Cut
slivers off the
top of the nail.
It helps the
quick recede.

HOW TO ~~MAKE YOUR~~ INDUCE VOMITING IN YOUR DOG

CALL YOUR VETERINARIAN OR ASPCA'S POISON HOTLINE AT
888-426-4435

FILL A SYRINGE OR TURKEY BASTER WITH THE BELOW AMOUNT OF 3%
HYDROGEN PEROXIDE. SQUIRT THE SOLUTION INTO THE BACK OF YOUR
DOG'S THROAT.

	10 LB. DOG	20 LB. DOG	40 LB. DOG	60 LB. DOG	80 LB. DOG
1 ST	5 cc.	10 cc.	20 cc.	30 cc.	40 cc.
DOSE	(1 teaspoon)	(2 teaspoons)	(4 teaspoons)	(2 tablespoons)	(2 ¾ tablespoons)

GATHER CLEANING SUPPLIES (RUBBER GLOVES, PLASTIC TRASH BAG,
PAPER TOWELS)

TAKE YOUR DOG OUTSIDE, DO NOT LEAVE UNATTENDED
WAIT FOR 15 MINUTES.

IF AFTER 15 MINUTES YOUR DOG DOES NOT VOMIT ADMINISTER A 2ND DOSE.

2 ND	5 cc.	10 cc.	20 cc.	30 cc.	40 cc.
DOSE	(1 teaspoon)	(2 teaspoons)	(4 teaspoons)	(2 tablespoons)	(2 ¾ tablespoons)

WAIT FOR 15 MINUTES.

IF YOUR DOG DOES NOT VOMIT, SEE YOUR VETERINARIAN IMMEDIATELY.
IF YOUR DOG DOES VOMIT, WAIT ANOTHER 15 MINUTES TO ENSURE YOUR
DOG GETS IT ALL OUT BEFORE TAKING TO SEE YOUR VETERINARIAN.


WATCH FOR SIGNS OF YOUR PET OVERHEATING




SYMPTOMS OF OVERHEATING INCLUDE:

- HEAVY PANTING
- EXCESSIVE THIRST
- GLAZED EYES
- VOMITING AND BLOODY DIARRHEA
- BRIGHT OR DARK RED TONGUE, GUMS
- STAGGERING
- ELEVATED BODY TEMPERATURE (104°F AND UP)
- WEAKNESS, COLLAPSE
- INCREASED PULSE AND HEARTBEAT
- SEIZURES
- EXCESSIVE DROOLING
- UNCONSCIOUSNESS

Heatstroke is Serious. Know the Signs.



HOW TO TREAT (AND POSSIBLY SAVE!) OVERHEATED DOG



Watch your pooch for signs of overheating during the hot weather.

1

Move your pet to a cooler area immediately, preferably with a fan.

2

Take your pet's temperature using a rectal thermometer.

3

Put cool, wet towels on

- neck,
- armpits
- between legs
- wet ear flaps and paw pads

4

Give your dog fresh cool drinking water.

5

Transport your pet to your vet right away!

Common foods toxic to dogs



Alcohol



Onions



Grapes



Coffee



Nuts



Avocado



Chocolate



Sugar free gum



Garlic

15 POISONOUS PLANTS FOR PETS

© University of Illinois Extension





**PET
POISON
HELPLINE®**

24/7 Animal Poison Control Center

800-213-6680

EARLY SIGNS OF BLOAT

DROOLING

LICKING LIPS

UNCOMFORTABLE 'WORRIED'

ATTEMPT TO VOMIT

PROGRESSION

STRAINING TO DEFECATE

ATTEMPT TO VOMIT - PHLEGM

DISTENDED ABDOMEN – BASKETBALL PING

DEPRESSED

COLLAPSE

DEATH